Michigan Collaborative Standardization of Compounded Oral Liquids

Dosage Form:

Suspension

60 days

Prug name: Flecainide

Acetate

Concentration: 20 mg/mL **Shelf Life:**

Route: Oral **Storage:** Refrigerate or Room Temp

Volume: 120 mL **Auxiliary Labeling:** Shake well

Ingredients	QS	Quantity	Units
Flecainide acetate 100 mg		24	Tablets
Vehicle	Υ	120	mL

Directions:

- 1. Triturate flecainide acetate tablets to a fine powder in a mortar
- 2. Levigate with approximately 20 mL of vehicle to form a paste
- 3. Add additional vehicle via geometric portions almost to volume, mixing thoroughly after each addition
- 4. Rinse the mortar and pestle with base and pour into graduated cylinder
- 5. QS with vehicle to 120 mL
- 6. Transfer to glass amber bottle
- 7. Shake well

Notes:

 Vehicles for which stability has been demonstrated include equal mixtures of Ora-Sweet:Ora-Plus, Ora-Sweet SF:Ora-Plus or cherry syrup

References:

• Loyd AV Jr. Int J Pharmaceut Compd. 1997;(1):103.

Reviewed: March 2014; SC/ME

The information provided on this site is only to be used for educational and informational purposes only. It may not be relied upon to direct or provide care to a patient, or to prepare treatments for a patient. This material is not intended as a substitute for retaining and consulting appropriate medical and health care professionals, who remain solely responsible to provide and direct patient care. Be aware that the information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professionals or pharmacists, and may require governmental approvals. The owner and provider of this site disclaims and assumes no liability for any injury and/or damage to persons from use of any information contained on this site.





