Michigan Collaborative Standardization of Compounded Oral Liquids

Drug name:	Procarbazine	Dosage Form: Suspension		
Concentration:	10 mg/mL	Shelf Life:	7 days	
Route:	Oral	Storage:	room temperature	
Volume:	50 mL	Auxiliary Labeling:	shake well, protect from light	

Ingredients	QS	Quantity	Units
Procarbazine 50-mg capsule		10	capsule
Glycerin		2	mL
Strawberry syrup	Y	50	mL

Directions:

- 1. Empty the contents of ten 50-mg procarbazine capsules into a mortar.
- 2. Add 2 mL glycerin and mix to a thick uniform paste.
- 3. Add 10 mL strawberry syrup in incremental proportions; mix until uniform.
- 4. Transfer the mixture to an amber glass bottle and rinse mortar and pestle with small amounts of strawberry syrup; add rinses to the bottle in sufficient quantity to make 50 mL.
- 5. Label "shake well" and "protect from light".

References:

• Taketoma CK, Kraus DM, Hodding JH. Pediatric Dosage Handbook. 19th Ed. Lexi-Comp, Inc. Hudson, OH, 2012.

Reviewed: April 2013; BW/ME

The information provided on this site is only to be used for educational and informational purposes only. It may not be relied upon to direct or provide care to a patient, or to prepare treatments for a patient. This material is not intended as a substitute for retaining and consulting appropriate medical and health care professionals, who remain solely responsible to provide and direct patient care. Be aware that the information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professionals or pharmacists, and may require governmental approvals. The owner and provider of this site disclaims and assumes no liability for any injury and/or damage to persons from use of any information contained on this site.



